



Scan to know paper details and  
author's profile

# A Review - Study on Phytochemical and Neuropharmacological Effects of Cassia Tora Flowers

*Tarique Hussain, Dr. Abdul Hafiz, Vishwakarma Ravi, Dr. Deepak Kumar & Dr. Imran Kazmi*

*Glocal University-Delhi-Yamunotri*

## ABSTRACT

Cassia tora flowers family-Leguminosae/ Fabaceae has great medicinal properties and used to treat various diseases. Flowers racemes terminal, oblong to rounded corymbose sepals 5, petals 5, yellow, perfect stamens 6-7. Contains chemical compounds like: Anthraquinone, chrysoferol, emodin, rhein, euphol, basselin, iso-stearic, palmitic, behenic acid etc., all chemicals isolated from this plant. These drugs affect the nervous system and its functioning, within the brain. There are still no effective therapies. Neurological related problems are so common today, that approximately 18% population suffer from disorders each year.

**Keywords:** phytochemical and neuropharmacological effects, cassia tora.

**Classification:** DDC Code: 341.1 LCC Code: HA154

**Language:** English



London  
Journals Press

LJP Copyright ID: 925642  
Print ISSN: 2631-8490  
Online ISSN: 2631-8504

London Journal of Research in Science: Natural and Formal

Volume 22 | Issue 2 | Compilation 1.0







# A Review - Study on Phytochemical and Neuropharmacological Effects of Cassia Tora Flowers

Tarique Hussain<sup>α</sup>, Dr. Abdul Hafiz<sup>σ</sup>, Vishwakarma Ravi<sup>ρ</sup>, Dr. Deepak Kumar<sup>ω</sup>  
& Dr. Imran Kazmi<sup>✧</sup>

## ABSTRACT

*Cassia tora* flowers family-Leguminosae/Fabaceae has great medicinal properties and used to treat various diseases. Flowers racemes terminal, oblong to rounded corymbose sepals 5, petals 5, yellow, perfect stamens 6-7. Contains chemical compounds like: Anthraquinone, chrysophanol, emodine, rhein, euphol, baselol, iso-istearic, palmitic, behenic acid etc., all chemicals isolated from this plants drugs that effects the nervous system and its functioning, within the brain. there are still no effective therapies Neurological related problems are so common today, that approximately 18% population suffer from disorders each year.

These disorders produces serious health problems like behavioral / cognitive syndrome, sleep disorders, peripheral disorders, epilepsy, neurodegenerative disorders, Parkinsonism, neoplasm and many others. Parts of *Cassia tora* are used as an antifungal, antihelmintic, diuretic, expectorant, and laxative, treatment of glaucoma, hypertension, and treatment of skin diseases, ringworm and itch. Pharmacological and phytochemical studies of flowers part of *Cassia Tora* plants which have been done in different parts.

**Keywords:** phytochemical and neuropharmacological effects, cassia tora.

**Author α:** Glocal University-Delhi-Yamunotri Marg (State Highwat75) Mirzapur Pole, Dist-Saharanpur-247121, Up, India.

**σ:** Glocal University-Delhi-Yamunotri Marg (State Highwat75) Mirzapur Pole, Dist-Saharanpur- 247121, up, india.

**ρ:** Arya College of Pharmacy-Jaipur, Rajsthan- 302028, India.

**ω:** Jss College of Pharmacy-Ooty, Tamilnaidu-643001.

**✧:** King Abdul-Aziz University, Jeddah, Saudi Arabia.

## I. INTRODUCTION

Medicinal herbs constitute the corner stone of traditional medicinal practice worldwide. These herbs are relatively cheap and easily available.

These medicinal plants represent a great deal of untapped reservoir of drugs and the structural diversity of their component molecule makes a valuable source of novel lead compounds. The world health organization about 80% of living organism like: Human, developing countries almost exclusively on traditional medicines for first health care needs. The present review traditional medicines uses and recent studies on the active isolated compounds. The whole plants as well as specific parts like: Roots, Leaves, Seeds and Flowers have been widely used in Indian and south Asian medicine it is an annual monsoon weed prevalent in wastelands having many antimicrobial properties, have the problem of pollution and health hazards of conventional agro chemical in view deals with conducted to test its seeds extract against the common aerial fungus, soil fungus and root fungus in forest eco-systems.

Furthermore, about 80% of the world population is dependent on plant-based drugs (WHO, 1996). In Nigeria and most developing countries of the world, rural and urban dwellers, literate or illiterate rely heavily on herbal preparations for the treatment of various diseases despite the availability of orthodox medicine the alternate system of medicine like Ayurveda, Sidda, Unnani,

and other tribal folklore medicines have significantly contributed to the health care of population of India. Today, these systems are not only complementary but also competitive in the

treatment of various diseases. Initially the materials employed in these traditional medicines were almost botanical origin.

## II. MATERIALS AND METHODS

### MATERIALS

<p style="text-align: center;"><i>Drug:</i></p> <p style="text-align: center;">Ibuprofen Pentazocine Chlorpromazine Valproic acid, Diazepam</p> <p style="text-align: center;"><i>Plant:</i></p> <p style="text-align: center;">The fresh leaves of <i>Cassia tora</i></p>	<p style="text-align: center;"><i>Reagents:</i></p> <p style="text-align: center;">Benedict's reagent. Barfoed's reagent. Million's reagent Dragendroff's reagent. Hager's reagent Mayer's reagent. Wagner's reagent</p> <p style="text-align: center;"><i>Chemicals:</i></p> <p style="text-align: center;">Petroleum ether Chloroform Ethanol</p>	<p style="text-align: center;"><i>Instruments:</i></p> <ul style="list-style-type: none"> <li>● Eddy's hot plate- Medicraft Pvt. Ltd, Ambala A-10-042 Whole Board</li> <li>● Soxhlet apparatus- Biotechnics, India. Double distillatory- Infusil-India Pvt. Ltd. Bangalore.(Mark 2000 DDQ-X2)</li> </ul> <p>Electronic weighing balance- Citizen scale, India Oral feeding needle- Space Lab, Nasik</p>
--	---	---

### PLANT PROFILE:

**Botanical Name-** Cassia Obtusifolia

**Synonym-** Cassia tora, Senna Obtusifolia

**Family-** Leguminosae

**Vernacular name-** Panwar, Penwaad , Sanjsaboya, Sangsaboya, Taarutaa, Ringworm plants, foetidecassia etc.

**Common (Indian) names:**

**Hindi:** Charota, Chakwad, Chakavat.

**Bengali & Oriya:** Chakunda

**Guajarati:** Kawaria

**Canarese:** Gandutogache

**Malayalam:** Chakramandrakam, takara

**Marathi:** Takala



### III. METHODOLOGY

The flowers of *Cassia tora* are collected and shade dried. Coarse powder is made from these dried flowers and subjected to extraction in increasing polarities. Various extracts are prepared by using suitable solvents like petroleum ether, chloroform, alcohol and aqueous solvent. Each crude extract obtained after evaporating the solvent subjected to preliminary phytochemical screening and these extracts are utilized for neuropharmacological activity. Stock solution of formulation under study was prepared freshly, on the day of experimentation using distilled water.

The dose of formulation-500mg/kg, was selected for the study, the selected doses represent 1/4 of 2000mg/kg. Healthy, Albino Swiss mice- the chosen experimental animals were maintained in our animal house (12:12 dark: light cycle), with adequate ventilation, hygienic conditions maintained on normal palliated diet and water ad libitum. A group of six animals were housed in polypropylene cage of 26 X 19 X 13cm on paddy husk bed and covered with stainless steel wire mesh 28 X 20.5 cm with provision for water and feed. Healthy, Albino Swiss mice, of either sex (unless otherwise specified) weighing between 18-25 gm. were employed for study. All experiments were performed in research lab all parameters of different tests were observed and recorded by person blind to treatment protocol.

For all experiments, group refers to group of six animals (n=6). Data generated from various experimental procedures were analyzed for statistical significance followed by Dennett's multiple comparison tests.

**Successive Solvent Extraction:** The powdered material was subjected to batch extraction in Soxhlet apparatus. The solvents used were petroleum ether, chloroform, alcohol and distilled water. The powdered material of *Cassia tora* leaves were evenly packed in a Soxhlet extractor for extraction for about 36 hours with different solvents.

**Preliminary photochemical investigation of extract:** Qualitative chemical tests were

conducted for Petroleum ether, ethanolic, Chloroform and aqueous extracts of flowers of *Cassia tora* to identify the various phytoconstituents. The phytochemical investigation showed presence of flavonoids, saponins, carbohydrates, starch, gum, proteins, tannin and phenolic compounds.

**Tests for Carbohydrates-** Molisch's test (General test)

**For Reducing Sugars-** Fehling's test, Benedict's test

**Test for Monosaccharide's-** Barfoed's test

**Tests for Non-Reducing Sugars-** 1. Test solution do not give response to Fehling's and Benedict's test. 2. Tannic acid test for starch: With 20% tannic acid, test solution was observed for precipitate

**Tests for Proteins-** Biuret test, Millon's test, Xanthoprotein test, Test for protein containing sulphur, Precipitation test,

**Tests for Steroids-** Salkowski Reaction, Liebermann-Burchard Reaction

**Tests for Amino Acids-** Ninhydrin test, Test for Tyrosine, Test for tryptophan

**Tests for Flavonoids-** Shinoda test, Ferric chloride test

**Tests for Alkaloids-** Dragendroff's test, Mayer's test, Hager's test, Wagner's test

**Tests for Tannins and Phenolic Compounds-** To 2-3 ml test solution, added few drops of following solutions and was looked for respective colouration or precipitate

**5% Ferric chloride solution:-** Deep blue-black coloured.

- 1) **Lead acetate solution:** - White precipitate.
- 2) **Gelatin solution:** - White precipitate.
- 3) **Bromine water:-** Decoloration of bromine water.
- 4) **Acetic acid solution:-** Red colour solution.
- 5) **Potassium dichromate:-** Red precipitate.

**Tests for Vitamins-** Test for Vitamin A, Test for vitamin C (Ascorbic acid), Test for Vitamin D

### IV. TESTS FOR GLYCOSIDES

**General test for Glycosides-Part A:** To 2-3 ml of extract dil H<sub>2</sub>SO<sub>4</sub> was added and heated on a

water bath for 1-2mins. Neutralise with 10% NaOH, check with litmus paper and to resulting solution add Fehling's A and B. Increased red precipitate in this case shows glycosides are present. Part B: To 2-3 ml of extract, water was added and heated. According to need, NaOH was added for neutralisation and also added equal quantity of water. To the resulting solution added Fehling's A and. Increased red precipitate in this case showed glycosides are absent.

*Tests for Cardiac Glycosides-* Baljet's test, Legal's test (For cardenoloids), Test for deoxysugars (Kellar Killani test), Libermann's test (For bufadenolids)

*Tests for Saponin Glycosides-* Foam test, Haemolytic test

## V. PHARMACOLOGICAL INVESTIGATION

ACUTE (ORAL) TOXICITY STUDY (Acute oral toxicity Fixed Dose Procedure, FDP)

Acute oral toxicity study for the proprietary formulation was carried out using OECD/OCED guideline 420 (modified adopted 23rd march 2006). The test procedure minimizes the number of animals required to estimate the oral acute toxicity of a chemical and in addition estimation of LD<sub>50</sub>, confidence intervals. The test also allows the observation of signs of toxicity and

can also be used to identify chemicals that are likely to have low toxicity.

*Principle of the FDP-* The fixed dose procedure is method for assessing acute oral toxicity that involve the identification of a dose level that cause evidence of non-lethal toxicity (termed evident toxicity) rather than a dose level that cause lethality. Evident toxicity is a term describing clear signs of toxicity following administration of test substance, such that an increase to the next highest fixed dose would result in the development of severe toxic signs and probably mortality.

*Procedure:* Healthy, young adult Albino Swiss mice (18-25gm), nulliparous and non-pregnant were used for this study. Food, but not water was withheld for 3-4 hours and further 1-2 hours post administration of sample under study. Fixed dose level of 5, 50, 500 mg/kg were initially chosen as dose level that would be expected to allow the identification of dose producing evident toxicity. During the validation procedure, a fixed dose of 2000mg/kg was added to provide more information on substance of low acute toxicity.

*Observation:* Animals were observed individually at least every 5 minutes once during first 30 minutes after dosing, periodically at 2hrs during the first 24 hours (with special attention during the first four hours) and daily thereafter, for a total of 14 days

## VI. RESULTS

Percentage Yield, Colour, Consistency and Solubility in Water of Different Extracts

Plant Part Used	Extracts	Percentage Yield	Color	Consistency	Solubility in Water
Cassia tora leaves	Petroleum	2.08	Dark green	Sticky	Insoluble
	Chloroform	1.95	Dark green	Sticky	Soluble
	Ethanol	1.96	Dark green	Sticky	Highly soluble
	Aqueous	2.13	Dark brown	Dry powder	Highly soluble

PRELIMINARY PHYTOCHEMICAL SCREENING of *Cassia tora*

Phytochemical constituents	Pet ether	Chloroform	Ethanol	Aqueous
Alkaloids	-	+	+	+
Glycosides	-	++	++	++
Carbohydrates	-	+	+	+

Flavonoids	-	+	+++	+++
Saponins	-	+	+++	+++
Tannins	-	+	+++	+++
Steroids	-	-	++	++
Proteins	+	+	+	+
Fats and oils	++	-	-	-
Starch	-	-	++	++
Gums	-	-	+++	+++
Phenolic compounds	-	-	+++	+++

- absent  
 + indicates  
 ++more clarity  
 +++ better response

### NEUROPHARMACOLOGICAL STUDY

*Effect of EEC and AEC on Exploratory behavior in mice:*


S.NO	Groups	Dose(mg/kg)	No.of Head dip
1	Control	-	19.2±1.12
2	Chlorpromazine	1	6.33±0.66 <sup>c</sup>
3	EEC	500	9.17±7.45 <sup>a,d</sup>
4	AEC	500	10.83±1.95 <sup>a,d</sup>

All values are mean±SEM, (n=6), <sup>a</sup>P<0.05, <sup>c</sup>P<0.001, when compared with control. <sup>d</sup>P<0.05 when compared with standard.

Effect of EEC and AEC flowers on exploratory behavior in mice (Percentage inhibition)

Sr. No.	Groups	Dose(mg/kg)	No, of head dip	Percentage Inhibition
1	Control	-	19.2±1.12	-
2	Standard (Chlorpromazine)	1	6.33±0.66 <sup>c</sup>	67.03-52
3	Ethanollic extract	500	9.17±7.45 <sup>a,d</sup>	52.23
4	Aqueous extract	500	10.83±1.95 <sup>a,d</sup>	43.59

*Effect of EEC and AEC leaves on exploratory behavior in mice:*




*Effect of EEC and AEC on diazepam induced sleeping time:*


Sr.No.	Groups	Dose(mg/kg)	Onset of Action (min)	Duration of Action (min)
1	Control(Diazepam)	5	121.8±5.5	83.00±2.04
2	EEC+Diazepam	500	83.83±2.15 <sup>b</sup>	114.7±3.48 <sup>b</sup>
3	AEC+Diazepam	500	54.00±1.65 <sup>b</sup>	126.0±1.92 <sup>b</sup>

all values are mean±SEM, n=6, <sup>b</sup>P<0.001, when compared with control

*Effect of EEC and AEC on diazepam induced sleeping time:*



*Effect of EEC and AEC on Maximal Electroshock (MES) Method Graph:*



**VII. CONCLUSION**

The present work suggests that it requires isolating and characterizing the active components responsible for the neuropharmacological activities and further studies to

reveal the exact mechanisms of action responsible for the neuropharmacological activities of Cassia tora flowers. Both the extracts EEC and AEC at the dose 500mg/kg body weight along with valproic acid at the dose 300mg/kg body weight showed significant reduction in time to recover

from electrically induced convulsions in mice when compared with control group ( $15.52 \pm 0.22, 18.33 \pm 0.16$  and  $28.17 \pm 1.01$ ,  $P < 0.001$ ,  $P < 0.001$  and  $P < 0.001$ ). The sedative-hypnotic effect of EEC and AEC was assessed using diazepam induced sleeping time. The preliminary phytochemical evaluation of different extracts of cassia tora flowers revealed the presence of flavonoids, carbohydrates, starch, gum, proteins, tannins, saponins and phenolic compounds. The results of phytochemical analysis were significant in ethanolic and aqueous extracts when compared with petroleum ether and chloroform extracts. In the present study effect of EEC and AEC on various parameters like exploratory behavior, analgesic activity, sleeping time and anti-convulsion activity was studied to expose the neuropharmacological property in different animal models.

## REFERENCES

- Dabriyal, RM, Narayana, DBA, Ayurvedic Herbal Raw Material, The Eastern Pharmacist, 1998, 31-35.
- Nadkarni, R.M., Indian Materia Medica, Vol I, Popular Book Depot, Mumbai, 1954, p.291.
- Shibata, S., Morishita, E., Kaheda, M., Kimura, Y., Takido, M and Takashashi, S., Pharm.Bull, 1969, 17,454.
- Green MM, Singer JM, Sutherland DJ, Hibben CR. 1991. Larvicidal activity of *Tagetes minuta* (marigold) toward *Aedes aegypti*. *J Am Mosq Control Assoc* 7:282–286.
- Jang YS, Baek BR, Yang YC, Kim MK, Lee HS. 2002. Larvicidal activity of leguminous seeds and grains against *Aedes aegypti* and *Culex pipiens pallens*. *J Am Mosq Control Assoc* 18:210–213. [13]. Jang, Y. S.; Baek, B. R.; Yang, Y. C.; Kim, M. K.; Lee, H. S. Larvicidal activity of leguminous seeds and grains against *Aedes aegypti* and *Culex pipiens pallens* (Diptera: Culicidae). *J. Am. Mosq. Control Assoc.* 2002, 18, 210-213.
- Zadikoff CM. 1979. Toxic encephalopathy associated with use of insect repellent. *J Paediatr* 95:140–142 [32]. WHO Malaria factsheet, 2014
- Yen, G. C.; Chung, D. Y. Antioxidant effects of extracts from *Cassia tora* L. Prepared under different degrees of roasting on the oxidative damage to biomolecules. *J. Agric. Food Chem.* 1999, 47, 1326-1332.
- WHO [World Health Organization]. 2002. The world health report 2002—reducing risks, promoting healthy life Geneva, Switzerland: World Health Organization [27].
- Wink M. 1993. Production and application of phytochemicals from an agricultural perspective. In: van Beek TA, Breteler H, eds. *Phytochemistry and agriculture* Oxford, United Kingdom: Clarendon Press. p 171–213.
- Choi, J. S.; Lee, H. J.; Park, K. Y.; Ha, J. O.; Kang, S. S. In vitro antimutagenic effects of anthraquinone aglycones and naphthopyrone glycosides from *Cassia tora*. *Planta Med.* 1997, 63, 11-14.
- Jain S, Patil K. Phytochemical and pharmacological profile of *Cassia tora* Linn-an overview. *Indian J Nat Pro Res.* 2010; 1(4): 430-7.
- Froestl W, Pfeifer A, Muhs A. Cognitive enhancers (Nootropic) Part 3 Drug interacting with targets other than receptor or enzymes disease modifying drugs. *J Alz Dis.* 2013; 34(1): 1-114.
- Pitchaimani V, et al. Nootropic activity of acetaminophen against Colchicin induced cognitive impairment in rats. *J Clin Biochem Nutr.* 2012; 50(3): 241-4.
- Ingkaninan K, Temkitthawon P, Chuenchom K. Screening for acetylcholinesterase inhibitory activity in plants used in Thai traditional rejuvenating and neurotonic remedies. *J Ethnopharmacol.* 2003; 89(2): 261-4.
- Murray A, Faraoni M, Castro M. Natural AChE inhibitors from plants and their contribution to Alzheimer's disease therapy. *Current Neuropharmacology* 2013; 11(4): 388-413. 18. Agnati LF, Genedani S, Leo G.  $\beta$  peptides
- Akbar S.M. Tariq and M. Nisa. A Study on CNS depressant activity of *Salvia haematodes* Wall. *Pharmaceutical Biology* 1984; 22(1): 41-44.
- Varma RK, Garg B.D. and Ahmad A. Pharmacodynamic studies on *Kalanchoe integrifolia*-an indigenous plant 1986; 18: 78-83.

18. Syed Kamil M, Liyakat Ahmed MD and Paramjyothi S. Neuropharmacological effects of ethanolic extract of *Portulaca quadrifida* Linn. In mice. *International journal of pharm Tech research* 2010; 2(2):1386-1390
19. Trease G.E and Evans M.C. *Textbook of Pharmacognosy*, 14th ed. London.2002; 81-90: 269-275,300.[http://en.wikipedia.org/wiki/Yh\\_e\\_Neurologist](http://en.wikipedia.org/wiki/Yh_e_Neurologist).
20. Ganeshchandra Sonavane, Vikram Sarveiya, Veena Kasture, Sanjay B. Kasture. Behavioral actions of *Myristica fragrans* seeds. *Indian Journal of Pharmacology* 2001; 33: 417-424
21. Radhakrishnan R., Zakaria M.N.M., Islam M.W. Chem H.B., Kamil M., Chan K and Al-Attas A. Neuropharmacological actions of *Portulaca oleraceae* L.V.Sativa (Hawk). *J. of Ethanopharm.* 200; 76: 171-176.
22. Shafiuddin MD., Liyakhat Ahmed MD., Taranalli A.D. and Khaja pasha. Influence of cyclohexanoyl thiosemicarbazide and some anticonvulsant drugs on neurotransmitter levels in rat-brain. *Int.J. of Chem.Sci.* 2009; 7(1): 264-272.
23. Ganeshchandra Sonavane, Vikram Sarveiya, Veena Kasture, Sanjay B. Kasture. Behavioral actions of *Myristica fragrans* seeds. *Indian Journal of Pharmacology* 2001; 33: 417-424.
24. Umar Kyari Sandabe, Patrick Azubuike Onyeyili, and Gregory Anene Chibuz. Sedative and anticonvulsant effects of aqueous extract of *Ficus sycomorus* L. (Moraceae) stem bark in rats. *Veterinarski Arhiv* 2003; 73 (2):103–110.
25. Tripathi KD. *Essential Medical pharmacology*, 2nd ed. Jaypee Brothers Publication, New Delhi.1988: 390-91.
26. Rahimi R., Nikfar S. and Abdullahi M. Efficacy and tolerability of *Hypericum perforatum* in major depressive disorder in comparison with selective serotonin reuptake inhibitors: A Metaanalysis. *Prog Neuropsychopharmacol. Biol. Psychiatry* 2009; 33:118-27.
27. Kokate CK., Purohit AP.,and Gokhale SB. *Textbook of Pharmacognosy*, 36th ed; Nirali Publication, Pune.2006:36,126.
28. Everitt B. J and Robbins T. W. Neural systems of reinforcement for drug addiction from actions to habits to compulsion. *Nature Neuroscience* 2005; 8(11): 1481-1489.
29. Hossein Hosseinzadeh and Vahid Khosravan. Anticonvulsant effects of aqueous and ethanolic extracts of *Crocus sativus* L. stigmas in mice. *Arch. Irn. Med.* 2002; 5(1):44-47.
30. Vision 2030, Project Directorate on Poultry. (2011) Available from: [http://www.pdonpoultry.org/pdpnew/images/extfiles/reports/PDP\\_Vision\\_2030.pdf](http://www.pdonpoultry.org/pdpnew/images/extfiles/reports/PDP_Vision_2030.pdf). Accessed on 27-02-2016.
31. McNamee, S.E., Cunningham, R. and Elliott, C.T. (2013).
32. Simultaneous immunochemical detection of four banned antibiotic growth promoters in raw and cooked poultry tissue. *Food Addit. Contam. A.*, 30: 1270-1278.